ALASKA JEPARTMENT OF FISH AND GI E 1/5-32-032

LAKE SURVEY SUMMARY

[16-31-10230-2067-3001-0010 Chilkat Lake
1.62 360 IQ' 34" (\$ \$3
MAP REF. Skagway B-3 Alaska 1 29 S R 56 E S 14 LAT. 59° 22' N LONG. 135° 56' W
OC. 3 miles SSW of Klukwan, near Haines, Alaska SURF. ELEV. 175' (Approx.)
1. SURF. ACRES 2,491 (DOT Est.) MAX. DEPTH 1851 AVG. DEPTH AVG. DEPTH ACRE FT. ACRE FT. SHOAL AREA Concentrated on WATER COLOR SECCHI READING
AQUATIC VEGETATION Very abundant higher plants on NW-SE shores of lake; submergent and emergent vegetation both present. Not identified - water lilly, emergent grasses seen
2. FISH SPECIES: NATIVE Ct, DV, RS, SS, Whitefish, Stickleback INTRODUCED None
3. FISHING HISTORY Reported to be the best Ct fishing in the Haines area. Fish 18-20 inches are relatively common. Average size reported to be near 16 inches. DV fishing is excellent through ice in winter but Ct are poor then.
4. INVERTEBRATES Aquatic
5. INLETS 7 major inlets were noted on map DISCHARGE cfs Due to rough lake conditions, survey was limited DISCHARGE cfs to lake - supplementary sheet will be used to DISCHARGE cfs
6. OUTLETS one - Colon Creek Estimated DISCHARGE 20-30 cfs 20-30 CFS DISCHARGE cfs BARRIERS None
7. SPAWNING AREAS None noted during this survey - Commercial Fish Division should have a good idea on this. Later observation will be put on supplementary sheet.
8. WATERSHED TYPE Land on SW side of lake is steep mtns. DRAINAGE AREA 38 square miles ACRES Land on NE side is rolling hills surrounded by heavy spruce forest. Also alder and 9. ACCESSIBILITY Accessible by river boat (jet or air), wheel plane or float plane. cottonwood.
10. ACCESS STATUS Public; there are many private summer cabins in choice locations along shore.
11. USE SHE One private hunting and fishing Lodge for public rental. Mostly private use.
12. OTHER USE Comm fisheries red salmon counting weir on outlet.
13. POLLUTION none noted
14. REMARKS Acres and Elev. from Tom Wade - State Div. of Lands - HNS. Depth, drainage, discharge from USFWS-River Basin Progress Report II-1961 Weather - cloudy - windy - rough water. Chilkat River Basin Study - USFWS 1959-61 should be consulted for comparisons and additional info. Water quality analysis was done for dissolved 02 and Ph, water taken at beginning of lake outlet. Readings were: D.O. 12 mg/l; Ph of 8. Water temp. was 61° F. A gill net - minnow trap, fish sampling summary was conducted. Attached supplementary sheet shows map of work done.

Chilket Lake

APPrix 59° 20' N. hat 138° 55' W. Ling. 3000 some I take type is gladier occurted take, Otear, against Mountain

2 Elevation 25 ft approx

3 Surrounding Cover is some timber, birch, cottonwood, alder, who

4 Flow Though The lake Max 1200 HJsm, Min 35 HJsm Norm. 100 HJsm

5 outlet length is approx. to milo, Slough-like - no ritles

6 No battier other Then Klohini + Chilket tiver mud,

7 Inlets - all from the mountoins and is snow rugoff.

8 Water Temp. Surface & 63°F

9 Oxygen Content -

10 Ph factor -

Il Fish are spowning in lake with none in wallet.

12 Species in lake - Cotthroat, Sticklebooks, Sockeye, Pinks Cham, Coho, occasional King

13 Plenting Record - nine

14 Food organius Visible - Sticklebooks, juscets rlorene

16 Ago groups present -

18 Paresites + diseases noted - only intestinal + contourie

17 Stomack contents - entirely stickle books

18 Remorks - Chilket lake is 8 min by air from Horines.

There is no established Trail To it nor shelter ofter one gets There. One has To Crass The Chilket R which is quite a barrier for books. It is 3 mi. from The Chilket R. To The lake. Dean Goodwin has a fish comp on one of the Islands in The lake. Flyin-ded.